

FORECASTING OF RAINFALL USING STATISTICAL DOWNSCALING
MODEL (SDSM) – GENERAL CIRCULATION MODEL (GCM) FOR FUTURE
ESTIMATION OF RAINWATER HARVESTING

‘AAINAA HATIN BINTI AHMAD TARMIZI

A thesis submitted in fulfillment of the requirement award of the
Master's Degree of Civil Engineering

Faculty of Civil Engineering and Built Environment
Universiti Tun Hussein Onn Malaysia

SEPTEMBER 2020

For my parents, who supported me with love and understanding. And also my sisters and my partner, whom has provided patience advice and guidance throughout the research process. Thank you all for your unwavering support.



PTTA UTHM
PERPUSTAKAAN TUNKU TUN AMINAH

ACKNOWLEDGEMENT

First and foremost, praises and thanks to the God, the Almighty, for His showers of blessings throughout my research work to complete the research successfully.

I would like to express deep and sincere gratitude to my main research supervisor, Dr Siti Nazahiyah Binti Rahmat for giving me the opportunity to do research and providing invaluable guidance throughout this research. Her dynamism, vision, sincerity and motivation have deeply inspired me. She has taught me the methodology to carry out the research and to present the research as clearly as possible. It was a great privilege and honor to work and study under her guidance. I am extending my heartfelt thanks to her late husband, Dr Ahmad Zurisman Bin Mohd Ali and her family for their acceptance and patience during meeting and discussion I had with her on my research journey. I would like to also thank my co-supervisor, Dr Nurul Nadrah Aqilah Binti Tukimat from Universiti Malaysia Pahang for the guidance especially on every steps of my journey learning climate modelling. Both of my supervisors definitely inspire me to become a successful woman in education in the future. I am extremely grateful for what they have offered me.

I am extremely grateful to my parents, Ahmad Tarmizi Bin Abd Karim and Kamaliah Binti Abdullah, for their love, prayers, caring and sacrifices for educating and preparing me for my future. Also, I express my thanks to my sisters, Auni Aqilah, Ainul Hayat and Amni Bahiah, for their support and valuable prayers. My special thanks goes to my partner, Muhammad Aiman Ramle for his love, understanding, prayers and continuing support to complete this research work.

ABSTRACT

Changes in the spatial and temporal rainfall pattern affected by the climate change need to be investigated as its significant characteristics are often used for managing water resources. In this study, the impacts of climate change on rainfall variability in Johor was investigated by using General Circulation Model (GCM) on the availability of daily simulation for three representative concentration pathways (RCP) scenarios, RCP2.6, RCP4.5 and RCP8.5 for interval year of $\Delta 2030$, $\Delta 2050$ and $\Delta 2080$. In addition, the annual future rainfall trend and harvested rainwater volume estimation for the first interval year of $\Delta 2030$ were also made. Daily rainfall series from eight (8) stations in Johor capturing 30 years period (1988-2017) with less than 10% missing data were chosen. Of all 26 predictors, only five (5) were chosen for each station to form a rainfall equation at each station for prediction analyses. It can be observed that the temperature (nceptemp), surface specific humidity (ncepshum) and near-surface relative humidity (nceprhum) had a strongest influence in the local weather formations with R values ranged from 0.5 to 0.7. The annual mean rainfall for RCP 2.6, 4.5 and 8.5 was predicted increase by of 17.5%, 18.1% and 18.3%, respectively as compared to historical data. Kluang was predicted to receive the highest amount of rainfall, and the lowest was in Segamat. Moreover, the Mann-Kendall test was used to detect the trend and resulted in no trend for RCP 2.6. Even so, RCP 4.5 showed a significant upward trend in Muar and Kota Tinggi, and for RCP 8.5, all regions were detected to have an upwards trend except for Pontian and Kluang. Volume harvested rainwater (V_R) was calculated, which resulted in increment of 16.8% for both RCP 2.6 and RCP 4.5. However, RCP 8.5 gave the highest possibility to store harvested rainwater with 18.3% increment in the future. In general, the concentration of greenhouse gases from RCP 8.5 gave the highest rainfall in future. Thus, the results will be useful in evaluating the performances of the rainwater harvesting system in Johor incorporating climate change for future research.

ABSTRAK

Perubahan corak hujan dalam masa dan ruang yang dipengaruhi oleh perubahan iklim perlu diselidiki kerana ciri-ciri pentingnya sering digunakan untuk mengurus sumber air. Dalam kajian ini, kesan perubahan iklim terhadap kebolehubahan curahan hujan di Johor disiasat dengan menggunakan *General Circulation Model* (GCM) terhadap ketersediaan simulasi harian untuk tiga *Representative Concentration Pathways*, RCP2.6, RCP4.5 dan RCP8.5 tahun selang $\Delta 2030$, $\Delta 2050$ dan $\Delta 2080$. Corak tahunan curahan hujan masa hadapan dan anggaran jumlah air hujan yang dituai untuk tahun selang pertama $\Delta 2030$ juga dibuat. Data hujan harian dari lapan (8) stesen di Johor yang merangkumi tempoh 30 tahun (1988-2017) dengan kurang dari 10% data yang hilang telah dipilih. Dari semua 26 ramalan, hanya lima (5) yang dipilih untuk setiap stesen untuk membentuk persamaan hujan di setiap stesen untuk analisis ramalan. Kajian ini menunjukkan bahawa suhu (nceptemp), kelembapan spesifik permukaan (ncepshum) dan kelembapan relatif permukaan (nceprhum) mempunyai pengaruh paling kuat dalam pembentukan cuaca setempat dengan nilai R berkisar antara 0.5 hingga 0.7. Hujan purata tahunan untuk RCP 2.6, 4.5 dan 8.5 diramalkan meningkat masing-masing 17.5%, 18.1% dan 18.3% berbanding dengan data sejarah. Kluang dijangka akan menerima jumlah curah hujan tertinggi, manakala Segamat terendah. Seterusnya, ujian Mann-Kendall digunakan untuk mengesan corak hujan dan hasil daripada analisis, RCP 2.6 tidak menunjukkan sebarang corak perubahan di masa hadapan. Walau bagaimanapun, RCP 4.5 menunjukkan corak kenaikan yang signifikan di Muar dan Kota Tinggi, dan untuk RCP 8.5, semua wilayah dikesan mengalami kecenderungan meningkat kecuali untuk Pontian dan Kluang. Isi padu takungan air hujan di hitung (V_R), akan mengalami kenaikan 16.8% untuk RCP 2.6 dan RCP 4.5. Walau bagaimanapun, RCP 8.5 berkemungkinan boleh menyimpan air hujan yang dituai dengan kenaikan 18.3% pada masa akan datang. Secara umum, kepekatan gas rumah hijau dari RCP 8.5 memberikan curahan hujan tertinggi pada masa depan. Oleh itu, hasilnya akan berguna dalam menilai prestasi sistem penuaian

air hujan di Johor yang menggabungkan perubahan iklim untuk penyelidikan masa hadapan.



TABLE OF CONTENT

TITLE	
DECLARATION	ii
DEDICATION	iii
ACKNOWLEDGEMENT	iv
ABSTRACT	v
ABSTRAK	vi
TABLE OF CONTENT	viii
LIST OF TABLES	xi
LIST OF FIGURES	xiii
LIST OF SYMBOL AND ABBREVIATIONS	xv
LIST OF APPENDICES	xvii
CHAPTER 1 INTRODUCTION	1
1.1 Background of study	1
1.2 Problem statement	3
1.3 Research objectives	4
1.4 Scope of research	4
1.5 Research Significance	6
1.6 Outline of the Thesis	7
CHAPTER 2 LITERATURE REVIEW	8
2.1 Introduction	8
2.2 Changes in rainfall with climate change	9

2.3	Review of climate models	12
2.3.1	Box model	13
2.3.2	Zero-dimensional model	14
2.3.3	Radiative-convective model	15
2.3.4	Energy balanced model	16
2.3.5	General circulation model	17
2.4	Downscaling as a supporting tool for General Circulation Model	22
2.4.1	Dynamical downscaling	23
2.4.2	Statistical downscaling	25
2.5	Trend Analysis of Rainfall	28
2.5.1	Mann-Kendall	28
2.5.2	Spearman's ratio	29
2.6	Rainwater harvesting (RWH) on simulated climate data	29
2.7	Summary	34
CHAPTER 3 RESEARCH METHODOLOGY		35
3.1	Introduction	35
3.2	Study Area	37
3.3	Missing Data	40
3.3.1	Arithmetic mean method	40
3.3.2	Normal ratio method	40
3.4	Statistical Downscaling Model (SDSM)	41
3.5	Retrieving climate predictors data	42
3.6	Screening of climate predictors	44
3.7	Scenario Generator on Projection of Future Rainfall	46
3.8	Mann-Kendall (MK) Test for Trend Analysis Test	48
3.9	Volume of harvested rainwater	50
CHAPTER 4 RESULTS AND DISCUSSION		51
4.1	Introduction	51
4.2	Climate Predictors Selection	52

	x
4.2.1 Calibration process	54
4.2.2 Validation process	55
4.2.3 Climate predictors	57
4.3 Projection of Future Rainfall using Canadian Earth System Model 2 (CanESM2)	59
4.4 Trend Analysis of Annual Projected Rainfall for all Stations	67
4.5 Rainwater harvesting potential under climate change	71
CHAPTER 5 CONCLUSIONS AND RECOMMENDATIONS	79
5.1 Conclusions of the study	79
5.2 Recommendations	81
REFERENCES	82
APPENDIX A	92
APPENDIX B	97
APPENDIX C	100



LIST OF TABLES

2.1	Previous studies on climate change impacts on rainfall pattern	11
2.2	General Circulation Models (GCMs) versions	19
2.3	26 predictors filenames and the corresponding variable (Wilby and Dawson, 2013)	20
2.4	Advantages and disadvantages dynamical downscale	24
2.5	Studies using RCM on dynamical downscaling method	24
2.6	Advantages and disadvantages of statistical downscaling	26
2.7	Previous studies using SDSM	27
2.8	Policies and guidelines of RWH in Malaysia	30
2.9	Implementation of the RWH system	32
3.1	Description of rainfall stations	39
4.1	Significant predictors from NCEP for rainfall	53
4.2	Statistical performance of rainfall modelling during calibration	54
4.3	Standard error (%) for validation	57
4.4	Predictor variables selected for predicting daily precipitation	58
4.5	Mean future rainfall in Johor	62
4.6	Z statistic values from the MK test for all stations	70
4.7	Estimation of harvested rainwater volume (m ³) using historical data (1988-2010)	74

4.8	Estimation of harvested rainwater volume (m ³) using RCP 2.6	75
4.9	Estimation of harvested rainwater volume (m ³) using RCP 4.5	76
4.10	Estimation of harvested rainwater volume (m ³) using RCP 8.5	77



LIST OF FIGURES

2.1	Diagram of energy sources and sinks of radiation contributing to climate (Sillmann <i>et al.</i> , 2017)	10
2.2	Observed and projected changes in annual average precipitation in Asia (IPCC, 2014)	11
2.3	Concept of climate modelling	13
2.4	The 'two-box' model (Loren, 2010)	14
2.5	EBMs components (Arrow indicate the direction of influence) (Shell and Somerville, 2005)	16
2.6	Conceptual structure of GCMs (Trzaska and Schnarr, 2014)	18
2.7	Land-sea mask in percentage (left) and orography/surface altitude in meters (right) for CanESM2 on the global Gaussian grid (64 lat x 128 long) (Government of Canada, 2019)	20
2.8	The concept of the spatial downscaling process (Trzaska and Schnarr, 2014)	23
3.1	Methodology Chart	36
3.2	Rainfall stations selected for the study	38
3.3	Steps on downscaling NCEP predictors and GCM predictors	42
3.4	Climate predictor for historical observation	43
3.5	Climate data for future observation	43
3.6	Process of screening NCEP climate predictors using SDSM	45

3.7	Schematic diagram of downscaling future rainfall data	47
3.8	Scenario generator function	48
4.1	Results of validation for all selected rainfall stations	56
4.2	Rainfall stations with climate predictors	58
4.3	Spatial distribution of historical rainfall data (1988 - 2017)	59
4.4	Percentage of projected change in average annual rainfall compared to average annual historical rainfall under RCP 2.6, RCP 4.5 and RCP 8.5 scenarios	63
4.5	Spatial distribution of RCP 2.6	64
4.6	Spatial distribution of RCP 4.5	65
4.7	Spatial distribution of RCP 8.5	66
4.8	Trend analysis for annual rainfall data of RCP 2.6	68
4.9	Trend analysis of annual rainfall for all stations	71
4.10	Total harvested rainwater comparison of historical data versus 3 RCPs ($\Delta 2030$)	78



LIST OF SYMBOL AND ABBREVIATIONS

P_x	-	Estimates the missing value
SE	-	Standard error
ABS	-	Absolute
Z	-	The standard test statistic
V_R	-	Volume harvested rainwater
R	-	mean monthly rainfall
H_{RA}	-	consider 10 household roof area (100 m^2)
R_c	-	runoff coefficient (0.75)
(R)	-	coefficient of correlation
GCMs	-	General Circulation Models
SDSM	-	Statistical Downscaling Model
RCPs	-	Representative Concentration Pathways
RWH	-	Rainwater harvesting
DID	-	Department of Irrigation and Drainage
NCEP	-	National Centres of Environmental Prediction
CanESM2	-	Canadian Earth System Model 2
RCE	-	Radiative-moist-convective equilibrium
EBMs	-	Energy Balanced Model (EBMs)
CanESM2	-	Canadian Earth System Model (CanESM2)
CTEM	-	Terrestrial Carbon Model
CMOC	-	Ocean Carbon Model
CCCma	-	The Canadian Centre For Climate Research Modelling and Analysis
WCRP	-	World Climate Research Program

IPCC	-	Intergovernmental Panel on Climate Change
RCM	-	Regional climate model
NEM	-	North-east monsoon
SWM	-	South-west monsoon
MK	-	Mann-Kendall test
WMO	-	World Meteorologist Organization
SR	-	Spearman's ratio



LIST OF APPENDICES

A	The process of selecting NCEP climate predictors using Statistical Downscaling Model (SDSM) 4.2.9	92
B	The process of downscaling future prediction rainfall of General Circulation Model (GCM) using Statistical Downscaling Model (SDSM) 4.2.9	97
C	Trend analysis of annual rainfall data for RCP 4.5 and RCP 8.5	100



PTTA UTHM
PERPUSTAKAAN TUNKU TUN AMINAH

CHAPTER 1

INTRODUCTION

1.1 Background of study

It is now widely accepted that global climate change results in a rise within the frequency and intensity of climatic extremes, such as droughts and floods (Dore, 2005). More intense rainfall events had occurred all over the world and Malaysia has also received the impacts of the changes. In November 2019, the south region of Malaysia had an intense rainfall for almost a week that led to flash flood and was one of the worst tragedies that have ever happened, especially in Johor as the number of flood victims rise rapidly (Shah *et al.*, 2019). Prediction of rainfall events and climate changes in the future will lead to preparation before extreme events occur.

Climate prediction models and statistical analysis are a great tool to estimate future rainfall data. Various type of climate prediction models can be found to predict any climate data such as rainfall, temperature, wind velocity and others from different studies (Thorpe *et al.*, 2001; Claussen *et al.*, 2002; Hock & Holmgren, 2005; Gagnon *et al.*, 2005; Robinson & Catling, 2012; Bajracharya *et al.*, 2018). However, the General Circulation Model (GCM) has been used in many studies all over the world to predict future rainfall, temperature and other climate properties. GCMs is the

infamously mathematically computer models that represent various global climate system of physical processes (Wilby, Dawson & Barrow, 2002).

GCM is a system of the many grid cells that represent horizontal and vertical areas on the Earth's surface which consider the potential level of greenhouse gases (GHGs). It computes water vapour and atmospheric cloud interactions, direct and indirect effects of aerosols on radiation and precipitation, changes in snow cover and sea ice, the storage of warmth in soils and oceans, surfaces fluxes of warmth and moisture, and large-scale transport of warmth and water by the atmosphere and oceans (Ghil & Robertson, 2000).

Canadian Earth System Model 2 (CanESM2) is one of the many center groups provide the GCMs data. Nevertheless, currently GCMs are depending on the Intergovernmental Panel on Climate Change (IPCC) Fifth Assessment Report (AR5) which that is the modification from the AR4 series in terms of expected %CO₂ and the climate variables. Proportionately, CanESM2 is an appropriate centre group to provide the GCMs data as CanESM2 represents a part of the modelling community's contribution to the IPCC Fifth Assessment Report (AR5) (Scinocca et al., 2008; IPCC, 2014b). IPCC AR5 made a finding of a new set of scenarios that provide time-dependent projections of atmospheric greenhouse gas (GHG) concentrations. The new scenarios are called Representative Concentration Pathways (RCPs) which consists four types of pathways; RCP 8.5, RCP 6, RCP 4.5 and RCP 2.6. Each number for each RCPs represents their forcing (Wayne, 2013). The RCPs are the latest iteration of the scenario process and it is important for climate change research to use RCPs data to investigate the plausible trajectories of different aspects of the future.

A statistical process is required to create a bridge between GCM scale and native scale because the GCM scale is just too vast. In this case, the downscaling method was chosen to minimize the GCM scale to the local scale. A process termed downscaling has been developed to formulate climate predictions at the local scale. Downscaling methods might be broadly classified into two categories, namely dynamical and statistical downscaling. Statistical is the most generally used compared to dynamical because it is often implemented easily in any region and cheap. In order to choose the suitable downscaling method, the desired spatial and temporal resolution of the climate information, resources and time constraints need to be considered (Wilby & Dawson, 2013).

Rainwater harvesting (RWH) system is known for capturing rainwater for local use. It has been implemented in all over countries in the world since years ago, and it has many different techniques and methods on implementing the system (Lo & Koralegedara, 2015; Alamdari *et al.*, 2018; Zhang *et al.*, 2019). RWH can also solve the water scarcity problem and decrease natural disaster phenomena. Water demand has been increasing worldwide at a rapid pace resulting in increasing demand caused by demographic changes, socioeconomic factors, changes in agricultural practices and climatic variation (Mohammed *et al.*, 2007). Hofman & Paalman (2014) concluded that rainwater harvesting could effectively reduced 13.9%, 30.2% and 57.7% of runoff volume in three cases of the maximum daily rainfall (207.2 mm), the average annual maximum daily rainfall (95.5 mm) and the critical rainfall of rainstorm (50 mm), respectively.

For most developing countries, RWH is listed together as the precise adaptation strategies that the water sector must undertake to deal with future global climate change, particularly in areas with high water demand. Some studies were administered to research the impact of global climate change on rainwater harvesting (RWH) supported by future rainfall variation (Fewkes, 2012; Kahinda *et al.*, 2010; Pina *et al.*, 2009; Helmreich & Horn, 2009).

1.2 Problem statement

Johor has been receiving increasing number of heavy rainfalls over the past years. According to the research made by Shafie (2009) a storm that blows from South East China Ocean and West Pacific Ocean cause heavy rainfall events that lead to major floods in December 2006 and January 2007. Kota Tinggi district was hit badly from the storms that brought 287 mm and 338 mm of rain in four (4) days recorded in Bandar Kota Tinggi for the year 2006 and 2007 respectively. There has been evidence made by the scientific community at NASA by Buis (2020) that as global temperatures increase, extreme precipitation will very likely increase as well but these researches have yet to have any significant effects on climate change in future rainfall intensities at a specific area of Johor.

Climate changes have affected rainfall intensities, especially in monsoon seasons which resulted in floods to occur without warning. Consequently, the rising temperature will lead to the unpredicted dry season. Besides, the rapid growth of population and expansion in urbanization may contribute to natural disasters. Besides, they have led to imposing growing demand and pressure on water resources (Che-Ani *et al.*, 2009). Rainwater harvesting is one of the ways to avoid water crisis as there will be increasing demand in future. Thus, it is important to estimate future changes in rainfall for finding suitable measures to mitigate this natural disaster problem but there has been little work exploring on the effect of climate change towards the amount of rainfall that could be harvested for a specific 100 m^2 of rooftop area in Johor.

1.3 Research objectives

The objectives of this study are;

- 1) To generate the long-term rainfall trend with respect to changes to the climate under three (3) different Representative Concentration Pathways Scenarios (RCPs).
- 2) To examine the changes in the trend of rainfall for interval year of $\Delta 2030$, $\Delta 2050$ and $\Delta 2080$.
- 3) To analyse the impact of climate change on the implementation of rainwater harvesting (RWH).

1.4 Scope of research

In order to achieve all the research objectives, the following tasks were carried out;

1. Collection of historical rainfall data (1988-2017) from the Department of Irrigation and Drainage (DID), Malaysia.

2. Projection of future rainfall for interval year of $\Delta 2030$, $\Delta 2050$ and $\Delta 2080$ using General Circulation Model with the climate variable data were retrieved from The Canadian Earth System Model 2 (CanESM2).
3. The climate modelling data were downscaled using Statistical Downscaling Model (SDSM).
4. Analyzing trend analysis for future rainfall data using the Mann-Kendall test.
5. Estimating amount of rainfall that could be harvested for a specific 100 m^2 of rooftop area for each selected station using Volume Harvested Formula.

The Canadian Earth System Model 2 (CanESM2) have 26 climate predictors. However, in this research validation of climate predictor selection for each stations was made and these are the climate modelling parameter used in this research; mean sea level pressure (ncepmslp), surface specific humidity (ncepshum), near surface relative humidity (nceprhum), mean temperature at 2m (nceptemp), relative humidity at 500hPa (ncepr500), relative humidity at 850hPa (ncepr850), 500hPa geopotential height (ncepp500) and 850hPa geopotential height (ncepp850).

There are three (3) major limitations in this study that could be addressed in future research. Firstly, eight (8) rainfall stations across Johor were chosen based on the data received from DID, Malaysia. The selected station represents each district in Johor that had less than 10% missing data (<10). Unfortunately, none of Mersing district rainfall stations can be used due to the insufficient daily rainfall data. Secondly, the geographical element was not considered in choosing the rainfall stations. Most of the stations had more than 10% missing data ($>10\%$) which led to difficulty in finding the most appropriate method to be used. Boughton (2007) stated that any rainfall data can be discarded if the catchments in which there were so many periods of missing data that the period available for calibration and validation for projection was too short. Thirdly, even though rainfall is associated well with temperature, however, temperature was not considered in the current study.

1.5 Research Significance

In this subchapter, the importance of the study and the potential end-results are deliberated generally as below:

- As was mentioned in Section 1.1, climate change has affected rainfall distribution across the world, including Malaysia. The latest flood event in early December 2019 that affected Johor proved that climate change gave impact to the rainfall variability, even though Johor did expect to receive the North-east monsoon where rain falls more than the usual month. However, the intensity of the rainfall was higher compared to the past years. Hence, the climate model needs to be implemented as there has been limited research on the prediction of rainfall events, specifically in Johor using the General Circulation Models (GCMs). This would help in projecting future rainfall for the next 90 years.
- A climate model is often too broad and difficult to understand. This study will explain the variety of the models but will highlight more on GCMs as the aim of this research will be using GCMs as the climate prediction model.
- Rainfall trend analysis assists in the identifications of possible trends in climatic parameters, including the wet and dry periods. No comprehensive research has been conducted on finding the trend for future rainfall events in Johor. Hence, a possible trend within the projected future rainfall are going to be determined.
- The estimation of harvested rainwater volume using future rainfall data in Johor could provide some information that can be used for future planning.

1.6 Outline of the Thesis

Chapter 1 explains the goals, the scopes of the study and therefore, the importance of the research. This chapter also provides an overall picture of the work activities carried out within the thesis.

Chapter 2 provides a thorough review of research associated with the global climate change impacts rainfall variability, climate modelling, a statistical tool to downscale climate data and rainwater harvesting implementation using projected rainfall data. A review paper on climate change effects on rainfall has been published in the International Journal of Integrated Engineering.

Chapter 3 provides the outline of the methods used, the study area, the rainfall data used and the climate predictors were chosen. The procedures of finding missing data of daily rainfall are also given. This chapter also presents preliminary analytical results on the assessment of climate predictors data that were downscaled to local scale using SDSM. The procedure of retrieving the climate predictors, screening of predictors, and calibration and validation processes to find the most appropriate climate predictors for each selected rainfall stations are also discussed.

Chapter 4 consists of main findings in this research which are the prediction of future rainfall in Johor, the future trend analysis and the results of the potential volume of rainwater that would be collected in the future. This chapter reveals the procedure of predicting rainfall using GCM which were downscaled using SDSM for three (3) different climate scenarios of GCMs predictors of Hadley Center General Circulation Model (CanESM2); RCP 2.6, RCP 4.5 and RCP 8.5. Besides, the future annual rainfall trend results are also presented. The developed equation of harvested rainwater volume is presented in this chapter and also the differences between past and future harvested rainwater were discussed.

Finally, Chapter 5 provides a summary and hence the principal inference is drawn from individual components of the analysis. Recommendations are also presented for future research.

CHAPTER 2

LITERATURE REVIEW

2.1 Introduction

This chapter reviews the impacts of climate change towards rainfall, the use of climate modelling to project future rainfall, supporting statistical tool to help lessen the scale of climate modelling data and RWH implementation. The changes to the climate is difficult to predict because of rapid growth of population, urbanization, industrialization, and irrigated agriculture that increases pressure on existing water resources and wish to be acknowledged by humankind (Lee *et al.*, 2016). Climate change will cause a general intensification of the earth's hydrological cycle within the next 100 years with generally increased in precipitation, evatranspiration, the occurrence of storms and significant changes in biogeochemical processes influencing water quality (Pandey *et al.*, 2003). In Malaysia, the awareness of climate change is still shallow. The public knew what the causes and effects of climate change are but tend to ignore the aftermath that could happen towards the earth.

Sea level rise is one of the significant impacts of climate change due to the global warming that causes ice from the south and north Antarctica to melt. Rainfall intensity was also affected by the changes in climate (Ahmad Tarmizi *et al.*, 2019). Rainfall varied from year to year and over decades and resulted in changes in the

REFERENCES

- Adefisan, E. (2018). Climate Change Impacts on Rainfall and Temperature Distributions Over West Africa from Three IPCC Scenarios. *Journal of Earth Science & Climate Change*, 9, 476.
- Ahmad Tarmizi, A. H. (2019). Climate Change and Its Impact on Rainfall. *International Journal of Integrated Engineering*, 11(1).
- Aladenola, O. O., & Adeboye, O. B. (2010). Assessing the potential for rainwater harvesting. *Water Resources Management*, 24(10), 2129-2137.
- Alamdari, N., Sample, D. J., Liu, J., & Ross, A. (2018). Assessing climate change impacts on the reliability of rainwater harvesting systems. *Resources, Conservation and Recycling*, 132, 178-189.
- Almazroui, M., Islam, M. N., Balkhair, K. S., Şen, Z., & Masood, A. (2017). Rainwater harvesting possibility under climate change: A basin-scale case study over the western province of Saudi Arabia. *Atmospheric research*, 189, 11-23.
- Al-Mukhtar, M., & Qasim, M. (2019). Future predictions of precipitation and temperature in Iraq using the statistical downscaling model. *Arabian Journal of Geosciences*, 12(2), 25.
- Amirabadizadeh, M., Ghazali, A. H., Huang, Y. F., & Wayayok, A. (2016). Downscaling daily precipitation and temperatures over the Langat River Basin in Malaysia: a comparison of two statistical downscaling approaches. *International Journal of Water Resources and Environmental Engineering*, 8(10), 120-136.
- Anandhi, A., Srinivas, V. V., Kumar, D. N., & Nanjundiah, R. S. (2009). Role of predictors in downscaling surface temperature to river basin in India for IPCC SRES scenarios using support vector machine. *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 29(4), 583-603.
- Ashok, K., & Yamagata, T. (2009). The El Niño with a difference. *Nature*, 461(7263), 481-484.
- Ayanlade, A., Radeny, M., Morton, J. F., & Muchaba, T. (2018). Rainfall variability and drought characteristics in two agro-climatic zones: An assessment of climate change challenges in Africa. *Science of the Total Environment*, 630, 728-737.
- Barros, V. R., & Field, C. B. (2014). *Climate change 2014: impacts, adaptation, and vulnerability. Part B: regional aspects*. Cambridge University Press.
- Barati, R., & Khazaei, S. (2018). Discussion of “Rainfall Trend Analysis of Hydrological Subbasins in Western Iran” by Mahsa Farhangi, Majid Kholghi, and Seyyed Ali Chavoshian. *Journal of Irrigation and Drainage Engineering*, 144(5), 07018012.

- Bajracharya, A. R., Bajracharya, S. R., Shrestha, A. B., & Maharjan, S. B. (2018). Climate change impact assessment on the hydrological regime of the Kaligandaki Basin, Nepal. *Science of the Total Environment*, 625, 837-848.
- Berezovskaya, S., Yang, D., & Kane, D. L. (2004). Compatibility analysis of precipitation and runoff trends over the large Siberian watersheds. *Geophysical Research Letters*, 31(21).
- binti Pg, D. S. N. A., Hasan, A., Ratnayake, U., Shams, S., Nayan, Z. B. H. & Rahman, E. K. A. (2018). Prediction of climate change in Brunei Darussalam using statistical downscaling model. *Theoretical and Applied Climatology*, 133(1-2), 343-360.
- Boulanger, J. P., Brasseur, G., Carril, A. F., De Castro, M., Degallier, N., Ereño, C., & Penalba, O. C. (2010). A Europe–South America network for climate change assessment and impact studies. *Climatic Change*, 98(3-4), 307-329.
- Boughton, W., & Chiew, F. (2007). Estimating runoff in ungauged catchments from rainfall, PET and the AWBM model. *Environmental Modelling & Software*, 22(4), 476-487.
- Buis, A. (2020, March 10). How Climate Change May Be Impacting Storms Over Earth's Tropical Oceans. <https://climate.nasa.gov/blog/2956/how-climate-change-may-be-impacting-storms-over-earths-tropical-oceans/>
- Bürger, G., Sobie, S. R., Cannon, A. J., Werner, A. T., & Murdock, T. Q. (2013). Downscaling extremes: An intercomparison of multiple methods for future climate. *Journal of Climate*, 26(10), 3429-3449.
- Campisano, A., Gnecco, I., Modica, C., & Palla, A. (2013). Designing domestic rainwater harvesting systems under different climatic regimes in Italy. *Water Science and Technology*, 67(11), 2511-2518.
- Che-Ani, A. I., Shaari, N., Sairi, A., Zain, M. F. M., & Tahir, M. M. (2009). Rainwater harvesting as an alternative water supply in the future. *European Journal of Scientific Research*, 34(1), 132-140.
- Christian Amos, C., Rahman, A., & Mwangi Gathenya, J. (2016). Economic analysis and feasibility of rainwater harvesting systems in urban and peri-urban environments: a review of the global situation with a special focus on Australia and Kenya. *Water*, 8(4), 149.
- Christensen, J. H. (2005). Prediction of regional scenarios and uncertainties for defining European climate change risks and effects (PRUDENCE). *Final Report, DMI*.
- Claussen, M., Mysak, L., Weaver, A., Crucifix, M., Fichet, T., Loutre, M. F., & Calov, R. (2002). Earth system models of intermediate complexity: closing the gap in the spectrum of climate system models. *Climate Dynamics*, 18(7), 579-586.
- Crosbie, R. S., Dawes, W. R., Charles, S. P., Mpelasoka, F. S., Aryal, S., Barron, O., & Summerell, G. K. (2011). Differences in future recharge estimates due to GCMs, downscaling methods and hydrological models. *Geophysical Research Letters*, 38(11).

- Danneberg, J. (2012). Changes in runoff time series in Thuringia, Germany–Mann-Kendall trend test and extreme value analysis. *Advances in Geosciences*, 31, 49-56.
- Darus, Z. M. (2009). Potential development of rainwater harvesting in Malaysia. *The 3rd WSEAS international conference on energy planning, energy saving, environmental education, Canary Islands, Spain*.
- Dore, M. H. (2005). Climate change and changes in global precipitation patterns: what do we know? *Environment International*, 31(8), 1167-1181.
- Eliseev, A. V., & Mokhov, I. I. (2007). Carbon cycle–climate feedback sensitivity to parameter changes of a zero-dimensional terrestrial carbon cycle scheme in a climate model of intermediate complexity. *Theoretical and applied climatology*, 89(1-2), 9-24.
- Emanuel, K., Wing, A. A., & Vincent, E. M. (2014). Radiative-convective instability. *Journal of Advances in Modelling Earth Systems*, 6(1), 75-90.
- Fewkes, A. (2012). A review of rainwater harvesting in the UK. *Structural Survey*, 30(2), 174-194.
- Feyissa, G., Zeleke, G., Bewket, W., & Gebremariam, E. (2018). Downscaling of future temperature and precipitation extremes in Addis Ababa under climate change. *Climate*, 6(3), 58.
- Fernandes, L. F. S., Terêncio, D. P., & Pacheco, F. A. (2015). Rainwater harvesting systems for low demanding applications. *Science of The Total Environment*, 529, 91-100.
- Field, C. B. (Ed.). (2014). *Climate change 2014–Impacts, adaptation and vulnerability: regional aspects*. Cambridge University Press.
- Flato, G. M., Boer, G. J., Lee, W. G., McFarlane, N. A., Ramsden, D., Reader, M. C., & Weaver, A. J. (2000). The Canadian Centre for Climate Modelling and Analysis global coupled model and its climate. *Climate Dynamics*, 16(6), 451-467.
- Fraedrich, K. (1978). Structural and stochastic analysis of a zero-dimensional climate system. *Quarterly Journal of the Royal Meteorological Society*, 104(440), 461-474.
- Gauthier, T. D. (2001). Detecting trends using Spearman's rank correlation coefficient. *Environmental Forensics*, 2(4), 359-362.
- Gagnon, S., Singh, B., Rousselle, J., & Roy, L. (2005). An application of the statistical downscaling model (SDSM) to simulate climatic data for streamflow modelling in Québec. *Canadian Water Resources Journal*, 30(4), 297-314.
- Gebremeskel, S., Liu, Y. B., De Smedt, F., Hoffmann, L., & Pfister, L. (2004). Analysing the effect of climate changes on streamflow using statistically downscaled GCM scenarios. *International Journal of River Basin Management*, 2(4), 271-280.
- GhaffarianHoseini, A., Tookey, J., GhaffarianHoseini, A., Yusoff, S. M., & Hassan, N. B. (2016). State of the art of rainwater harvesting systems towards promoting

green built environments: a review. *Desalination and Water Treatment*, 57(1), 95-104.

- Ghil, M. and Robertson, A. W. (2000). Chapter 10 Solving problems with GCMs: General circulation models and their role in the climate modelling hierarchy', *International Geophysics*, 70(C), 285–325.
- Ghisi, E., Bressan, D. L., & Martini, M. (2007). Rainwater tank capacity and potential for potable water savings by using rainwater in the residential sector of southeastern Brazil. *Building and Environment*, 42(4), 1654-1666.
- Ghisi, E., Montibeller, A., & Schmidt, R. W. (2006). Potential for potable water savings by using rainwater: An analysis of over 62 cities in southern Brazil. *Building and Environment*, 41(2), 204-210.
- Giorgi, F. (2005). Climate change prediction. *Climatic Change*, 73(3), 239-265.
- Government of Canada. (2019, April 18). The ensemble of daily predictor variables developed from the CanESM2 CMIP5 experiments. <http://climate-scenarios.canada.ca/index.php?page=canesm2-predictor-notes>
- Hansen, J., Kharecha, P., Sato, M., Masson-Delmotte, V., Ackerman, F., Beerling, D. J., & Rockstrom, J. (2013). Assessing “dangerous climate change”: required reduction of carbon emissions to protect young people, future generations and nature. *PloS one*, 8(12), e81648.
- Harvey, L. D. (1988). A semianalytic energy balance climate model with explicit sea ice and snow physics. *Journal of Climate*, 1(11), 1065-1085.
- Hasan, Z., Akhter, S., & Islam, M. (2014). Climate change and trend of rainfall in the south-east part of coastal Bangladesh. *European Scientific Journal*, 10(2).
- Hassan, Z., & Harun, S. (2012). Application of statistical downscaling model for long lead rainfall prediction in Kurau River catchment of Malaysia. *Malaysian Journal of Civil Engineering*, 24(1).
- Helmreich, B., & Horn, H. (2009). Opportunities in rainwater harvesting. *Desalination*, 248(1-3), 118-124.
- Herrmann, T., & Schmida, U. (2000). Rainwater utilisation in Germany: efficiency, dimensioning, hydraulic and environmental aspects. *Urban water*, 1(4), 307-316.
- Hessami, M., Gachon, P., Ouarda, T. B., & St-Hilaire, A. (2008). Automated regression-based statistical downscaling tool. *Environmental Modelling & Software*, 23(6), 813-834.
- Hewitt, C. D. (2004). Ensembles-based predictions of climate changes and their impacts. *Eos, Transactions American Geophysical Union*, 85(52), 566-566.
- Hock, R., & Holmgren, B. (2005). A distributed surface energy-balance model for complex topography and its application to Storglaciären, Sweden. *Journal of Glaciology*, 51(172), 25-36.
- Hofman, J. A. M. H., & Paalman, M. (2014). *Rainwater harvesting, a sustainable solution for urban climate adaptation?* KWR Watercycle Research Institute.
- Hughes, L., & Steffen, W. (2003). Climate change in Victoria: trends, predictions and

- impacts. *Proceedings of the Royal Society of Victoria*, 125(1), 5-13.
- Hussain, M., Yusof, K. W., Mustafa, M. R., & Afshar, N. R. (2015). Application of statistical downscaling model (SDSM) for long term prediction of rainfall in Sarawak, Malaysia. *Water Resources Management*, 8, 269.
- Imteaz, M. A., Ahsan, A., Naser, J., & Rahman, A. (2011). Reliability analysis of rainwater tanks in Melbourne using daily water balance model. *Resources, Conservation and Recycling*, 56(1), 80-86.
- IPCC, 2014 (2014) *CLIMATE CHANGE 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects, Psychological Science*.
- IPCC (2014a) *Climate Change 2014 Part A: Global and Sectoral Aspects, Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*.
- IPCC (2014b) 'Climate Change 2014 Synthesis Report', *IPCC Fifth Assessment Report*, p. 80.
- Jacob, D., Elizalde, A., Haensler, A., Hagemann, S., Kumar, P., Podzun, R., & Teichmann, C. (2012). Assessing the transferability of the regional climate model REMO to different coordinated regional climate downscaling experiment (CORDEX) regions. *Atmosphere*, 3(1), 181-199.
- Jing, X., Zhang, S., Zhang, J., Wang, Y., & Wang, Y. (2017). Assessing efficiency and economic viability of rainwater harvesting systems for meeting non-potable water demands in four climatic zones of China. *Resources, Conservation and Recycling*, 126, 74-85.
- Jing, X., Zhang, S., Zhang, J., Wang, Y., Wang, Y., & Yue, T. (2018). Analysis and modelling of stormwater volume control performance of rainwater harvesting systems in four climatic zones of China. *Water resources management*, 32(8), 2649-2664.
- Jupp, T. E., & Cox, P. M. (2010). MEP and planetary climates: insights from a two-box climate model containing atmospheric dynamics. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1545), 1355-1365.
- Kabiri, R., Bai, R. V., & Chan, A. (2013). Regional precipitation scenarios using a spatial statistical downscaling approach for Klang watershed, Malaysia. *Journal of Environmental Research and Development*, 8(1), 126.
- Karumuri, A. K., & Yamagata, T. (2009). Climate change: El Niño with a difference. *Nature*, 461, 481-484.
- Kadioğlu, M. (1997). Trends in surface air temperature data over Turkey. *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 17(5), 511-520.
- Kahinda, J. M., Taigbenu, A. E., & Boroto, R. J. (2010). Domestic rainwater harvesting as an adaptation measure to climate change in South Africa. *Physics and Chemistry of the Earth, Parts A/B/C*, 35(13-14), 742-751.

- Khairoutdinov, M., & Emanuel, K. (2013). Rotating radiative-convective equilibrium simulated by a cloud-resolving model. *Journal of Advances in Modelling Earth Systems*, 5(4), 816-825.
- Kisakye, V., Akurut, M., & Van der Bruggen, B. (2018). Effect of climate change on reliability of rainwater harvesting systems for Kabarole District, Uganda. *Water*, 10(1), 71.
- Koukidis, E. N., & Berg, A. A. (2009). Sensitivity of the Statistical DownScaling Model (SDSM) to reanalysis products. *Atmosphere-Ocean*, 47(1), 1-18.
- Lee, K. E., Mokhtar, M., Hanafiah, M. M., Halim, A. A., & Badusah, J. (2016). Rainwater harvesting as an alternative water resource in Malaysia: potential, policies and development. *Journal of Cleaner Production*, 126, 218-222.
- Litofsky, A. L., & Jennings, A. A. (2014). Evaluating rain barrel stormwater management effectiveness across climatology zones of the United States. *Journal of Environmental Engineering*, 140(4), 04014009.
- Lo, K., & Koralegedara, S. (2015). Effects of climate change on urban rainwater harvesting in Colombo city, Sri Lanka. *Environments*, 2(1), 105-124.
- Lorenz, R. D. (2010). The two-box model of climate: limitations and applications to planetary habitability and maximum entropy production studies. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1545), 1349-1354.
- Lorenz, R. D. (2001). Correction to "Titan, Mars and Earth: Entropy production by latitudinal heat transport". *Geophysical Research Letters*, 28(16), 3169-3169.
- Mahmood, R., & Babel, M. S. (2013). Evaluation of SDSM developed by annual and monthly sub-models for downscaling temperature and precipitation in the Jhelum basin, Pakistan and India. *Theoretical and Applied Climatology*, 113(1-2), 27-44.
- Mann, H. B. (1945). Nonparametric tests against trend. *Econometrica: Journal of the Econometric Society*, 245-259.
- Maraun, D., Wetterhall, F., Ireson, A. M., Chandler, R. E., Kendon, E. J., Widmann, M., & Venema, V. K. C. (2010). Precipitation downscaling under climate change: Recent developments to bridge the gap between dynamical models and the end user. *Reviews of Geophysics*, 48(3).
- Mohammed, T. A., Noor, M. J. M. M., & Ghazali, A. H. (2007). Study on potential uses of rainwater harvesting in urban areas. In *Proceedings of the Colloquium on Rainwater Utilisation, Putrajaya, Malaysia, 19&20 April*.
- Mondal, A., Kundu, S., & Mukhopadhyay, A. (2012). Rainfall trend analysis by Mann-Kendall test: A case study of the north-eastern part of Cuttack district, Orissa. *International Journal of Geology, Earth and Environmental Sciences*, 2(1), 70-78.
- Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., Van Vuuren, D. P., & Meehl, G. A. (2010). The next generation of scenarios for climate change research and assessment. *Nature*, 463(7282), 747.
- Mrad, D., Djebbar, Y., & Hammar, Y. (2018). Analysis of trend rainfall: Case of north-eastern Algeria. *Journal of water and land development*, 36(1), 105-115.

- Murphy, B. F., & Timbal, B. (2008). A review of recent climate variability and climate change in southeastern Australia. *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 28(7), 859-879.
- Musayev, S., Burgess, E., & Mellor, J. (2018). A global performance assessment of rainwater harvesting under climate change. *Resources, Conservation and Recycling*, 132, 62-70.
- Nadiga, B. T., & Urban, N. M. (2019). Improved representation of ocean heat content in energy balance models. *Climatic Change*, 152(3-4), 503-516.
- Pachauri, R. K., Allen, M. R., Barros, V. R., Broome, J., Cramer, W., Christ, R., & Dubash, N. K. (2014). Climate change 2014: synthesis report. Contribution of Working Groups I, II and III to the fifth assessment report of the Intergovernmental Panel on Climate Change, 151.
- Pina, C. L., Kassaye, R. B., & Schaldach, R. (2009). Rainwater Harvesting Methods.
- Pandey, D. N., Gupta, A. K., & Anderson, D. M. (2003). Rainwater harvesting as an adaptation to climate change. *Current Science*, 85(1), 46-59.
- Partal, T., & Kahya, E. (2006). Trend analysis in Turkish precipitation data. *Hydrological Processes: An International Journal*, 20(9), 2011-2026.
- Peñuelas, J., & Filella, I. (2009). Phenology feedbacks on climate change. *Science*, 324(5929), 887-888.
- Phan-Van, T., Nguyen-Xuan, T., Van Nguyen, H., Laux, P., Pham-Thanh, H., & Ngo-Duc, T. (2018). Evaluation of the NCEP Climate Forecast System and its downscaling for seasonal rainfall prediction over Vietnam. *Weather and Forecasting*, 33(3), 615-640.
- Quadros, L. E. D., Mello, E. L. D., Gomes, B. M., & Araujo, F. C. (2019). Rainfall trends for the State of Paraná: present and future climate. *Revista Ambiente & Água*, 14(2).
- Rahmat, S. N., Jayasuriya, N., & Bhuiyan, M. A. (2015). Precipitation trends in Victoria, Australia. *Journal of Water and Climate Change*, 6(2), 278-287.
- Rahmat, S. N., Al-Gheethi, A. A. S., Ayob, S., & Shahli, F. M. (2020). Development of dual water supply using rooftop rainwater harvesting and groundwater systems. *SN Applied Sciences*, 2(1), 85.
- Robinson, T. D., & Catling, D. C. (2012). An analytic radiative-convective model for planetary atmospheres. *The Astrophysical Journal*, 757(1), 104.
- Thenmozhi, M., & Kottiswaran, S. V. (2016). Analysis of rainfall trend using Mann-Kendall test and the Sen's slope estimator in Udumalpet of Tirupur district in Tamilnadu. *International Journal of Agricultural Science and Research*, 6(2), 131-138.
- Sa'adi, Z., Shahid, S., Chung, E. S., & bin Ismail, T. (2017). Projection of spatial and temporal changes of rainfall in Sarawak of Borneo Island using statistical downscaling of CMIP5 models. *Atmospheric research*, 197, 446-460.
- Sammathuria, M. K., & Ling, L. K. (2009, August). Regional Climate Observation and Simulation of Extreme Temperature and Precipitation Trends. In *14th*

International Rainwater Catchment Systems Conference (pp. 3-6).

- Schneider, S. H., & Dickinson, R. E. (1974). Climate modelling. *Reviews of Geophysics*, 12(3), 447-493.
- Scinocca, J. F., McFarlane, N. A., Lazare, M., Li, J., & Plummer, D. (2008). The CCCma third generation AGCM and its extension into the middle atmosphere. *Atmospheric Chemistry and Physics*, 8(23), 7055-7074.
- Sen, P. K. (1968). Estimates of the regression coefficient based on Kendall's tau. *Journal of the American statistical association*, 63(324), 1379-1389.
- Şen, Z., Al Alsheikh, A., Al-Turbak, A. S., Al-Bassam, A. M., & Al-Dakheel, A. M. (2013). Climate change impact and runoff harvesting in arid regions. *Arabian Journal of Geosciences*, 6(1), 287-295.
- Shell, K. M., & Somerville, R. C. (2005). A generalized energy balance climate model with parameterized dynamics and diabatic heating. *Journal of Climate*, 18(11), 1753-1772.
- Sillmann, J., Thorarinsdottir, T., Keenlyside, N., Schaller, N., Alexander, L. V., Hegerl, G., & Zwiers, F. W. (2017). Understanding, modelling, and predicting weather and climate extremes: Challenges and opportunities. *Weather and Climate Extremes*, 18, 65-74.
- Silva, C. M., Sousa, V., & Carvalho, N. V. (2015). Evaluation of rainwater harvesting in Portugal: Application to single-family residences. *Resources, Conservation and Recycling*, 94, 21-34.
- Silva, H. R., Bhardwaj, R., Phelan, P. E., Golden, J. S., & Grossman-Clarke, S. (2009). Development of a zero-dimensional mesoscale thermal model for urban climate. *Journal of Applied Meteorology and Climatology*, 48(3), 657-668.
- Singh, A., Acharya, N., Mohanty, U. C., & Mishra, G. (2013). Performance of multi model canonical correlation analysis (MMCCA) for prediction of Indian summer monsoon rainfall using GCMs output. *Comptes Rendus Geoscience*, 345(2), 62-72.
- Svirezhev, Y. M., & von Bloh, W. (1997). Climate, vegetation, and global carbon cycle: the simplest zero-dimensional model. *Ecological Modelling*, 101(1), 79-95.
- Shafie, A. (2009). A Case study on floods of 2006 and 2007 in Johor, Malaysia. *Colorado State University*.
- Shah, MF., Nordin, R., Then, S., Murali, RSN & Khoo D. (2019, December, 16). More and more evacuees: Number rise as floods now move southwards to Johor and Pahang. *THE STAR*, 3.
- Syafrina, A. H., Zalina, M. D., & Juneng, L. (2015). Historical trend of hourly extreme rainfall in Peninsular Malaysia. *Theoretical and Applied Climatology*, 120(1-2), 259-285.
- Tabari, H., Marofi, S., Amini, A., Talaei, P. H., & Mohammadi, K. (2011). Trend analysis of reference evapotranspiration in the western half of Iran. *Agricultural and Forest Meteorology*, 151(2), 128-136.

- Takahashi, K. (2009). Radiative constraints on the hydrological cycle in an idealized radiative–convective equilibrium model. *Journal of the Atmospheric Sciences*, 66(1), 77-91.
- Tarafder, R., & Nandy, D. (2017). A Data Driven, Zero-Dimensional Time Delay Model with Radiative Forcing for Simulating Global Climate. *arXiv preprint arXiv:1709.08860*.
- Thenmozhi, M., & Kottiswaran, S. V. (2016). Analysis of rainfall trend using Mann-Kendall test and the Sen's slope estimator in Udumalpet of Tirupur district in Tamil Nadu. *International journal of agricultural science and research*, 6(2), 131-138.
- Thorpe, R. B., Gregory, J. M., Johns, T. C., Wood, R. A., & Mitchell, J. F. B. (2001). Mechanisms determining the Atlantic thermohaline circulation response to greenhouse gas forcing in a non-flux-adjusted coupled climate model. *Journal of Climate*, 14(14), 3102-3116.
- Trenberth, K. E. (2011). Changes in precipitation with climate change. *Climate Research*, 47(1-2), 123-138.
- Trzaska, S., & Schnarr, E. (2014). A review of downscaling methods for climate change projections. *United States Agency for International Development by Tetra Tech ARD*, 1-42.
- Tukimat, N. N. A., Syukri, N. A., & Malek, M. A. (2019). Projection the long-term ungauged rainfall using integrated Statistical Downscaling Model and Geographic Information System (SDSM-GIS) model. *Heliyon*, 5(9), e02456.
- Van Oldenborgh, G. J., Reyes, F. D., Drijfhout, S. S., & Hawkins, E. (2013). Reliability of regional climate model trends. *Environmental Research Letters*, 8(1), 014055.
- Vautard, R. (2018). Regional Climate Projections. In *Weather & Climate Services for the Energy Industry* (pp. 139-149). Palgrave Macmillan, Cham.
- Wallace, C. D., Bailey, R. T., & Arabi, M. (2015). Rainwater catchment system design using simulated future climate data. *Journal of Hydrology*, 529, 1798-1809.
- Wayne, G. (2013). *The Beginner's Guide to Representative Concentration Pathways*. England, Devon: SkepticalScience.
- Wilby, R. L., Dawson, C. W., & Barrow, E. M. (2002). SDSM—a decision support tool for the assessment of regional climate change impacts. *Environmental Modelling & Software*, 17(2), 145-157.
- Wilby, R. L., Charles, S. P., Zorita, E., Timbal, B., Whetton, P., & Mearns, L. O. (2004). Guidelines for the use of climate scenarios developed from statistical downscaling methods. *Supporting material of the Intergovernmental Panel on Climate Change, available from the DDC of IPCC TGCIA*, 27.
- Wilby, R. L., & Dawson, C. W. (2013). The statistical downscaling model: insights from one decade of application. *International Journal of Climatology*, 33(7), 1707-1719.
- Wing, A. A., Reed, K. A., Satoh, M., Stevens, B., Bony, S., & Ohno, T. (2018).

Radiative-convective equilibrium model intercomparison project. *Geoscientific Model Development*, 11, 793-813.

Wong, C. L., Yusop, Z., & Ismail, T. (2018). Trend of daily rainfall and temperature in Peninsular Malaysia based on gridded data set. *International Journal*, 14(44), 65-72.

Xu, C. Y. (1999). From GCMs to river flow: a review of downscaling methods and hydrologic modelling approaches. *Progress in Physical Geography*, 23(2), 229-249.

Yue, S., Pilon, P., & Cavadias, G. (2002). Power of the Mann–Kendall and Spearman's rho tests for detecting monotonic trends in hydrological series. *Journal of Hydrology*, 259(1-4), 254-271.

Youn, S. G., Chung, E. S., Kang, W. G., & Sung, J. H. (2012). Probabilistic estimation of the storage capacity of a rainwater harvesting system considering climate change. *Resources, Conservation and Recycling*, 65, 136-144.

Zhang, S., Zhang, J., Yue, T., & Jing, X. (2019). Impacts of climate change on urban rainwater harvesting systems. *Science of The Total Environment*, 665, 262-274.

Zulkarnain, H., Mohd Faez, M. M., & Ain Nihla, K. (2019). Rainfall projection corresponding to climate scenarios based on Statistical Down-Scaling Model over Perlis, Malaysia. *Journal of Engineering Research and Education*, 11, 9-14



PTTA
PERPUSTAKAAN TUNKU TUN AMINAH